



# Cardiac Congenital Pathologies More Frequent in Children in a Hospital of Guayaquil (Ecuador)

Nancy Jácome Guerrero<sup>1</sup>, Arianna Soto Andrade<sup>1</sup>, Carlos Jácome Pilco<sup>2</sup>, Darwin Núñez Torres<sup>2\*</sup>

<sup>1</sup>Universidad Católica Santiago de Guayaquil, Av. Pdte. Carlos Julio Arosemena Tola, CP: 090615, Guayaquil, ECUADOR

<sup>2</sup>Universidad Estatal de Bolívar, Centro de Investigación y Desarrollo Biotecnológico, CP: 020150, Guaranda, ECUADOR

\*Corresponding Author: [isba@doctor.upv.es](mailto:isba@doctor.upv.es)

**Citation:** Jácome Guerrero N, Soto Andrade A, Jácome Pilco C, Núñez Torres D. Cardiac Congenital Pathologies More Frequent in Children in a Hospital of Guayaquil (Ecuador). *Electron J Gen Med.* 2020;17(5):em226. <https://doi.org/10.29333/ejgm/7889>

## ARTICLE INFO

Received: 25 Feb. 2020

Accepted: 16 Mar. 2020

## ABSTRACT

Congenital heart diseases (CHD) are structural malformations of the heart or large vessels, which cause a functional impairment, present at birth, being discovered at that time or during the course of the life of the carrier. CHD are a group of complex diseases that cause a serious public health problem. The objective of the work was, to identify the most frequent congenital heart malformations in children under 1 year of age in the general hospital of the north IESS Los Ceibos of the city of Guayaquil (Ecuador) in the period July 2017- January 2018. The data of this investigation were taken from pediatric patients by reviewing medical records. In the study period, 35,105 pediatric care visits were made to children under 1 year of age. Around 1.025 (2.92%) cases of congenital malformations were detected, of which 108 corresponded to congenital heart malformations (CHD), that is, 10.53% of the malformations observed were of the CHD type. The most frequent cardiac congenital malformations in children under 1 year old are non-cyanotic congenital cardiac malformations (NC) with a total of 102 cases (94%) and 6 cases (6%). they correspond to congenital cardiac malformations (C), being these less frequent.

**Keywords:** malformations, congenital heart disease, cyanosis, non-cyanosis, prevalence, children

## INTRODUCTION

Congenital heart diseases (CHD) are structural malformations of the heart or the great vessels, which cause a functional impairment, which are present at birth, being discovered at that moment or during the course of the life of the carrier. This type of pathologies is an alteration in the embryonic development of the heart (3rd and 10th week of gestation) (1).

The causes of CHD are unknown, although there is evidence that inheritance plays a decisive role in 8% of those affected and teratogens are involved in 2% of cases. The genesis of the remaining 90% is multifactorial (hereditary predisposition) (2).

The CHD are highlighted due to their high lethality, whose mortality in children under one year corresponds to something more than 1/3 worldwide (3). While in our country they correspond to one of the main causes of mortality child category within the group of "catastrophic diseases", occupying between the third and fourth place in this list (4).

The most prevalent CHD are those of the acyanotic type (non-cyanotic), among the most frequent are: defects of the interventricular septum (VSD) and atrial septal defect (ASD), patent ductus arteriosus, pulmonary valvular stenosis and coarctation of the aorta (1).

The most prevalent CHD are those of the acyanotic type (non-cyanotic), among the most frequent are: defects of the interventricular septum (VSD) and atrial septal defect (ASD),

patent ductus arteriosus, pulmonary valvular stenosis and coarctation of the aorta (5).

Approximately 3-4% of all newborns have a congenital malformation, either structural or functional. Among these anomalies are congenital heart diseases (CHD), which are among the leading causes of death in the population of childhood malformations in Latin America (6,7). It is proposed that out of every 1.000 live births, 8 to 14 will have some type of CHD, with premature children being the most at risk (8).

In Ecuador, in 2011 and 2012 the National Institute of Statistics and Census (INEC) reported that CHD are among the first ten etiologies of infant mortality (4).

Heart diseases are usually accompanied by extra cardiac abnormalities and the relationship between sexes is very even, except in specific pathologies such as the transposition of large arteries where males are mostly affected with major ratio (9).

The presentation of CHD can be of two basic forms, cyanotic or cyanotic, the method of choice for the diagnosis of these malformations in pediatric patients is echocardiography, considering the advantages in costs and easy access (11,12).

Usually the most frequent pathologies are the non-cyanotic heart diseases being the ventricular septal defect (VSD), the atrial septal defect (ASD) and the patent ductus arteriosus (PDA) the most frequent in this group (8).

## CHD not Cyanosantes

Among the heart diseases that produce volume overload. The common pathophysiological denominator of this group is

the communication between the systemic and pulmonary sides of the circulation. These are characterized by pulmonary congestion. The most frequent representatives of this group are: atrial septal defect, ventricular septal defect, septal defects and persistent ductus arteriosus (9,10).

Among the cardiopathies that give rise to pressure overloads. The common pathophysiological denominator is the obstruction to normal blood flow. The manifestations are: sweating, tiraje, tachypnea, galloping rhythm, murmurs or recurrent pneumonia, cardiomegaly. The most frequent are outflow tract obstructions (pulmonary valve stenosis, aortic valve stenosis, and coarctation of the aorta). Obstructions of the entrance tract (mitral or tricuspid stenosis) are less frequent (9).

### CHD Cyanantes

These CHDs present with cyanosis and hypoxemia as a capital sign. This group can be subdivided according to the pathophysiology: if the pulmonary blood flow is reduced or increased (10). Among the cyanotic cardiopathies with decreased pulmonary blood flow.

These heart diseases are characterized by both obstruction to the pulmonary flow, either at the level of the tricuspid valve, the right ventricle or the pulmonary valve. Common heart diseases in this group include tricuspid atresia, tetralogy of Fallot, and various forms of single ventricle with pulmonary stenosis (9,10).

### CHD Cyanosias More Frequent

#### CHD with decreased pulmonary flow

Tricuspid atresia. It is the anatomic absence of the tricuspid valve, accompanied by severe hypoplasia or absence of a right ventricle. It manifests with severe hypoxemia and acidosis, it is considered a bad prognosis heart disease when it coexists with other heart diseases such as transposition of large arteries or pulmonary stenosis (3,9).

Tetralogía de Fallot: It has 4 components; obstruction to the exit in right ventricle, ventricular septal defect, overload of the aorta and hypertrophy of the right ventricle (9,10).

### Most Frequent Accianantes CHD

*Ventricular Septal Defect (VSD)*. It is a lesion in which the interventricular septum allows communication between the two ventricles.

Interventricular communication is classified according to the anatomical location where the defect is found: Membranous in 80%, Trabecular in 0.5 to 20%, Entrance tract in 5 to 7%, outflow tract (or infundibular) in 5% (3).

The purpose of this study was to know which are the most common congenital heart diseases in children under one year of age in the General Hospital of the North IESS Los Ceibos during the period from June 2017 to January 2018, it should be noted that the technical-scientific literature has been revised on the subject, there are no published antecedents of the epidemiology of CHD in this hospital or in the city in recent years, and there are very few studies on this subject in Ecuador.

## METHODS

Children under 1 year of age diagnosed with congenital heart diseases were registered in the general hospital of the

North IESS Los Ceibos during the period from June 2017 to January 2018. For this, the population of 108 individuals was used; randomization and the previous review of the clinical histories was carried out.

After identifying the pediatric patients under 1 year of age treated in the general hospital of the north IESS. Los Ceibos during the analysis period diagnosed with congenital cardiac pathologies, the prevalence of congenital cyanotic and non-cyanotic cardiac pathologies was determined.

Distributing this percentage according to sex, gestational age, birth weight, APGAR at the first minute of the patients in the study and according to certain characteristics of the mother such as age, number of feets and infectious factor that may be present.

For all previously described, a descriptive, observational, retrospective, cross-sectional analysis was applied.

## RESULTS AND DISCUSSIONS

In our study of a total of 35.105 pediatric visits to children under 1 year of age in the IESS Los Ceibos General Hospital in the period of analysis, 1.025 cases of congenital malformations were detected, accounting for 2,92% of total pediatric care; of these cases, 108 corresponded to congenital heart malformations (CHD), that is, 10,53% of the malformations observed were of the CHD type.

The prevalence obtained was 3 cases per 1,000 children under 1 year (0,30%) seen in this hospital. With respect to gender, the prevalence of CHD in the female population is relatively greater with 57 cases (52%) while in the male gender there were 51 cases (48%).

In this regard, studies conducted by Tassinari *et al.* (6) and Egbe *et al.* (7), indicate that approximately 3-6% of all newborns have a significant congenital malformation at birth, whether structural or functional; This suggests that the results of this study to date show us that this rate obtained more than 10 years ago, still remains in this sector of the country.

Précoma *et al.* (3), point out that, among the group of congenital malformations, heart diseases are the most frequent, leading to an incidence of 4 to 12 per 1,000 live births, and they are second only to congenital malformations worldwide. Premature children are the most at risk of having a CHD (8), whose mortality in children under one year corresponds to something more than 1/3 worldwide.

Studies carried out in our country by the National Institute of Statistics and Censuses (4), during the years 2011 and 2012 report that CHD malformations are among the first ten etiologies of infant mortality in Ecuador cataloged within the group of "catastrophic diseases", occupying between the third and fourth place in this list (4).

Of the 108 cases determined as congenital heart malformations (CHD), congenital non-cyanotic cardiac malformations stand out in 94%, with a total of 102 cases and the remaining 6% (6 cases) correspond to congenital malformations of cyanotic heart (C) (**Figure 1**).

### CONGENITAL CARDIAC MALFORMATIONS




Figure 1. Congenital Cardiac Malformations

Of the 102 cases that correspond to CHD NC, the most frequent was the CIA with 58 cases (56.86%), followed by patent ductus arteriosus (PDA) (19; 18.6%), VSD (18; 17.64%). There were 2 cases (1.96%) with CIA + CIV, and with aortic coarctation (CoA). Finally, in 1 case (0.98%) the pathologies were aortic stenosis (AE), ventricular hypoplasia, and abnormal pulmonary venous drainage (APVD) (Figure 2).

Regarding the 6 cases that correspond to cyanotic congenital heart malformations, they can be classified into different pathologies: TF with 5 cases representing 83.33%, this being the most prevalent pathology; and, transposition of large vessels with 1 case, which represents 16.66 (Figure 3).

Table 1. Study variables

VARIABLE NAME	VARIABLE	TIPE	MEASUREMENT LEVEL
Congenital Cardiac Malformations	Categorical Variable	Nominal	Yes/No
Congenital Cardiac Malformations	Categorical Variable	Nominal	Yes/No
Congenital Cardiac Malformations	Categorical Variable	Nominal	Yes/No
Age	Numerical Variable	Continuous	1, 2, 3 months, etc...

For the study of Cardiac Congenital Pathologies, the variables selected for this research are divided into pediatric aspects related to gestational age, birth weight and APGAR at minute of life; and, maternal aspects that may have influence on the appearance of congenital heart malformations, which are: maternal age, number of feets and presence of infectious factor.

For the variable gestational age we find the following results:

- A. Severe prematurity Nb (Newborn) less than 32 SG 8 cases corresponding to 7.4% of the total of our study cases. In these cases, the following congenital cardiac pathologies were presented: PAC 5 case – 62.5%, CIA 2 cases - 25%, CIV 1 case – 12.5%.
- B. Moderately premature Nb from 32 SG to 34 SG 9 cases corresponding to 8.33% of the total of our case studies. In these, the following congenital cardiac pathologies were presented: CIA 5 cases – 55.6%, PCA 2 cases – 22.2%, CIV 1 case – 11.1%, TF 1 case – 11.1%.

### CONGENITAL CARDIAC MALFORMATIONS, NOT CYANOSANTS




Figure 2. Congenital Cardiac Malformations, Non-Cyanosine

### CONDITIONAL CONGENITAL HEART MALFORMATIONS




Figure 3. Congenital Cardiac Heart Malformations

C. Nb prematurely mild from SG 35 to SG 36 17 cases corresponding to 15.74% of the total of our case studies. In these, the following congenital cardiac pathologies were presented: CIA 7 cases – 41.1%, PDA 5 cases – 29.4%, CIV 2 cases – 11.8%, and CoA TDGV, CIA + CIV with 1 case – 5.9 % each.

D. Nb to term 73 cases corresponding to 67.59% of the total of our case studies. In these the following congenital heart pathologies were presented: CIA 43 cases – 58.9%, CIV 14 cases – 19.1%, PDA 8 cases – 10.9%, TF 3 cases – 4.1% and CoA, EA, DVPA, CIA + CIV, HV with 1 case – 1.4% each.

E. Post-term Nb 1 case corresponding to 0.92% of the total of our study cases. In these the following congenital cardiac pathologies CIV 1 case -100% were presented.

For the variable APGAR at minute we find the following results:

A) APGAR at the minute of 0 to 3.13 cases corresponding to 12.03% of the total of our case studies. In these, the following congenital cardiac pathologies were presented: CIA 7 – 53.8%, PCA 3 cases – 23.1%, CIV 2 cases – 15.4%, AD 1 case – 7.7%.

B) APGAR at the minute from 4 to 6,22 cases corresponding to 20-37% of the total of our case studies. The following congenital cardiac pathologies were present in these cases: 13 cases – 59.2%, CIV 5 cases – 22.7%, PCA 3 cases – 13.6%, TDGV 1 case – 4.5%

C) APGAR at the minute of 7 or greater, 73 cases corresponding to 67.60% of the total of our case studies. The following congenital cardiac pathologies were presented in these cases: 34 cases – 46.5%, CIV 17 cases – 23.3%, PCA 15 cases – 20.5%, TF 4 cases 5.6%, CoA 2 cases – 2.7%

## ACKNOWLEDGEMENTS

We thank the IEES Los Ceibos Hospital, which allowed us and gave us the facilities to carry out this research work and in a special way, Dr. Fuad Guaman Garaicoa, who guided us in this research, and our parents, and our universities for the support provided.

## REFERENCES

- Uribe Rivera A, Diaz-Velez C, Cerron-Rivera C. Características epidemiológicas y clínicas de las cardiopatías congénitas en menores de 5 años del Hospital Almanzor Aguinaga Asenjo: Enero - diciembre 2012. *Horiz. Med.*, Lima 2015;15(1). <https://doi.org/10.24265/horizmed.2015.v15n1.07>
- Valdés Silva Y, Sánchez Ramírez E, Fuentes S. Malformaciones congénitas relacionadas con los agentes teratógenos. *Correo Científico Médico*, 2018;22(4):652-66. Available at: [http://scielo.sld.cu/scielo.php?script=sci\\_arttext&pid=S1560-43812018000400011&lng=es](http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S1560-43812018000400011&lng=es)
- Précoma D.B, Oliveira Gláucia M, Simão A, Dutra O, Coelho O, Izar A, de Oliveira MC et al. Updated Cardiovascular Prevention Guideline of the Brazilian Society of Cardiology - 2019. *Arq. Bras. Cardiol*, 2019;113(4):787-891. <https://doi.org/10.5935/abc.20190204> PMID:31691761 PMCID:PMC7020870
- INEC, Instituto Nacional de Estadísticas y Censos. Anuario de Estadísticas Hospitalarias: Egresos y Camas 2014; pp 522. Available at: [http://www.ecuadorencifras.gob.ec/documentos/web-inec/Estadisticas\\_Sociales/Camas\\_Egresos\\_Hospitalarios/Publicaciones-Cam\\_Egre\\_Host/Anuario\\_Camas\\_Egresos\\_Hospitalarios\\_2014.pdf](http://www.ecuadorencifras.gob.ec/documentos/web-inec/Estadisticas_Sociales/Camas_Egresos_Hospitalarios/Publicaciones-Cam_Egre_Host/Anuario_Camas_Egresos_Hospitalarios_2014.pdf)
- Echeverría D, López L, Arellano M. Detección temprana de cardiopatías congénitas en neonatos a término mediante la saturación de oxígeno medida por oximetría de pulso en el Hospital gineco-obstétrico Isidro Ayora Quito, 2013. *Revista médica Universidad Central del Ecuador*; 2015.
- Tassinari S, Martínez-Vernaza S, Erazo-Morera N, Pinzón-Arciniegas M.C, Gracia G, Zarante I. Epidemiología de las cardiopatías congénitas en Bogotá, Colombia en el período comprendido entre 2001 y 2014: ¿Mejoría en la vigilancia o aumento en la prevalencia? *Biomédica*, 2018;38:141-8. <https://doi.org/10.7705/biomedica.v38i0.3381> PMID:29809331
- Egbe A, Uppu S, Lee S, Stroustrup A, Ho D, Srivastava S. Congenital Malformations in the Newborn Population: A Population Study and Analysis of the Effect of Sex and Prematurity. *Pediatrics and Neonatology*, 2015;56(1):25-30. <https://doi.org/10.1016/j.pedneo.2014.03.010> PMID:25267275
- Mendieta-Alcántara GG, Santiago-Alcántara E, Mendieta-Zerón H, et al. Incidencia de las cardiopatías congénitas y los factores asociados a la letalidad en niños nacidos en dos hospitales del Estado de México. *Gac Med Mex* 2013;149:617-23.
- Kimchi A. Abstracts, International Academy of Cardiology Annual Scientific Sessions 2017 22nd World Congress on Heart Disease Vancouver, BC, Canada, Cardiology, 2017; pp. 349. Available at: <https://www.karger.com/Article/Pdf/477751>
- Hernández-Benítez R, Becerra-Becerra R. Ecocardiografía funcional en cuidados intensivos neonatales: experiencia en un hospital de tercer nivel. *Bol. Med. Hosp. Infant. Mex.* 2016;73(5):325-30. <https://doi.org/10.1016/j.bmhmx.2016.07.003> PMID:29384125
- Arguello J, Sánchez L, Tapia M, Mosso M, Flores F, Guamán J, Muñoz M, Bayas-Morejón F. Factors that Affect Cesarean Delivers in the General Hospital of Guaranda (Ecuador). *Asian Journal of Pharmaceutics*, 2018;12(2):S636-S639. <https://doi.org/10.22377/ajp.v12i02.2408>
- Patra S, Rama Sastry U, Mahimaiha J, Subramanian A, Shankarappa R, Nanjappa M. Spectrum of cyanotic congenital heart disease diagnosed by echocardiographic evaluation in patients attending paediatric cardiology clinic of a tertiary cardiac care centre. *Cardiology in the Young*, 2015;25(5):861-7. <https://doi.org/10.1017/S1047951114000882> PMID:24914874